

US006905612B2

(12) United States Patent

Dorian et al.

(10) Patent No.: US 6,905,612 B2

(45) **Date of Patent:** Jun. 14, 2005

(54) PLASMA CONCENTRATE APPARATUS AND METHOD

(75) Inventors: Randel Dorian, San Diego, CA (US);

Richard Wood Storrs, San Francisco,

CA (US)

(73) Assignee: Hanuman LLC, San Francisco, CA

(US)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 259 days.

- (21) Appl. No.: 10/394,828
- (22) Filed: Mar. 21, 2003
- (65) **Prior Publication Data**

US 2004/0182788 A1 Sep. 23, 2004

- (51) **Int. Cl.**⁷ **B01D 37/00**; B01D 15/02; B01D 24/00; B01D 29/00

(56) References Cited

U.S. PATENT DOCUMENTS

3,850,369 A	11/1974	Bull et al.
3,897,343 A	7/1975	Ayers
3,909,419 A	9/1975	Ayres
3,931,018 A	1/1976	North, Jr.
3,982,691 A	9/1976	Schultz
4,046,699 A	9/1977	Zine, Jr.
4,055,501 A	10/1977	Cornell
4,077,396 A	3/1978	Wardlaw et al.
4,187,979 A	2/1980	Cullis et al.
4,322,298 A	3/1982	Persidsky
4,416,654 A	11/1983	Schoendorfer et al.

4 464 467 A	0/4004	61 16 11
4,464,167 A	8/1984	Schoendorfer et al.
4,675,117 A	6/1987	Neumann et al.
4,776,964 A	10/1988	Schoendorfer et al.
4,818,386 A	4/1989	Burns
5,019,243 A	5/1991	McEwen et al.
5,053,127 A	10/1991	Schoendorfer et al.
5,131,907 A	7/1992	Williams et al.
5,141,645 A	8/1992	Shiraki et al.
5,147,290 A	9/1992	Jonsson
5,165,938 A	11/1992	Knighton

(Continued)

FOREIGN PATENT DOCUMENTS

EP	1 079 224 A2	2/2001
WO	WO 98/12274	2/1998
WO	WO 99/67277	12/1999

Primary Examiner—John Kim (74) Attorney, Agent, or Firm—William B. Walker

(57) ABSTRACT

A plasma concentrator for producing plasma concentrate from a plasma from which erythrocytes have been substantially removed. The device comprises a concentrating chamber having an inlet port and an concentrate outlet, the concentrating chamber containing hydrogel beads and at least one inert agitator; and a concentrate chamber having an inlet communicating with the concentrator outlet through a filter, and having an plasma concentrate outlet port. A process for producing plasma concentrate from plasma from which erythrocytes have been substantially removed, comprising the steps of a) moving the plasma into a concentrating chamber containing hydrogel beads and an agitator to form a hydrogel bead-plasma mixture; b) causing the agitator to stir the hydrogel bead-plasma mixture, facilitating absorption of water by the beads from the plasma, until a hydrogel bead-plasma concentrate is formed; and c) separating the plasma concentrate from the hydrogel beads by passing the plasma concentrate through a filter. The concentrator can be one or more syringe devices coupled for multiple concentrations.

29 Claims, 6 Drawing Sheets

